

Mean platelet volume level in patients with generalized anxiety disorder

Behice Han Almis & Ihsan Aksoy

To cite this article: Behice Han Almis & Ihsan Aksoy (2018) Mean platelet volume level in patients with generalized anxiety disorder, Psychiatry and Clinical Psychopharmacology, 28:1, 43-47, DOI: [10.1080/24750573.2017.1385210](https://doi.org/10.1080/24750573.2017.1385210)

To link to this article: <https://doi.org/10.1080/24750573.2017.1385210>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 11 Oct 2017.



Submit your article to this journal [↗](#)



Article views: 37290



View related articles [↗](#)




View Crossmark data [↗](#)



Citing articles: 9 View citing articles [↗](#)

Mean platelet volume level in patients with generalized anxiety disorder

Behice Han Almis and Ihsan Aksoy 

Psychiatry Department, Adiyaman University Research and Education Hospital, Adiyaman, Turkey

ABSTRACT

OBJECTIVE: Generalized anxiety disorder (GAD) is characterized by persistent and chronic anxiety state. Stressful life events and anxiety usually cause an increase in platelet volume and activity through various mechanisms. Mean platelet volume (MPV), which is indicative of platelet size, is accepted as an indication of platelet activity.

METHOD: In this first retrospective cohort study about MPV among GAD patients, we compared complete blood count especially in terms of platelet count (PLT), platelet distribution width (PDW), plateletcrit (PCT) and MPV values of 60 GAD patients with 60 healthy controls.

RESULTS: MPV was found to be significantly higher ($p = .008$) and platelet count was found to be significantly lower in the GAD group ($p = .001$). The area under the ROC curve (AUC) of MPV levels for GAD was 0.655 (95% CI: 0.557–0.754, $p < .001$).

DISCUSSION: Increased MPV levels in GAD patients in our study supports the hypothesis of increased platelet activation due to sympathetic system activation. Also platelet levels were found to be significantly lower in GAD patients in our study. This finding is consistent with the non-linear inverse relation between platelet volume and platelet count. According to ROC curve analysis for differentiation of GAD from those in the control group, MPV would be considered to be a “poor” biochemical marker. We believe that further controlled prospective studies about this issue will be valuable.

ARTICLE HISTORY

Received 17 July 2017

Accepted 22 September 2017

KEYWORDS

Generalized anxiety disorder; complete blood count; mean platelet volume; platelet count; platelet distribution volume; plateletcrit

Introduction

Generalized anxiety disorder (GAD) is characterized by persistent and chronic anxiety state that lasts for at least 6 months which takes place in multiple areas of life. The lifetime prevalence is thought to be between 5.7% and 14.2% [1]. GAD patients are more likely to be encountered at primary health care facilities with non-specific complaints such as headache, muscle stiffness, gastrointestinal complaints, back pain and insomnia [2]. Major depressive disorder is a common companion to GAD beside increased rates of alcohol and substance abuse. It also aggravates physical diseases especially cardiovascular disorders [2,3].

Increased platelet activity has been detected in patients with major depression with a history of myocardial infarction and also in patients without any other risk factors effecting platelet activity [4,5]. On the other hand, anxiety symptoms in patients with coronary artery disease have been shown to be more effective in increased platelet activity than depression [6]. In a meta-analysis, Roest et al. [3] suggested that anxiety increased the risk of developing coronary artery disease by 26% when other medical conditions were controlled.

The exact mechanism of increased platelet activity after mental stress remains unclear. Anxiety disorders related “peripheral changes” are considered to be a

cumulative sign of the total central nervous system activity. Stressful life events and anxiety usually elevate circulating levels of blood catecholamines. Increased catecholamine levels are thought to increase thrombosis by causing an increase in platelet activity and sympathoadrenal activation is thought to stimulate platelets through the α -2 receptors, resulting in an increase in platelet volume and activity [7,8]. It is accepted that serotonin which plays an important role in anxiety disorders is also an important factor in determining vascular tone and platelet aggregation [9]. Plasma platelets have the ability of synthesis, release and reuptake of serotonin similar to neurons in the central nervous system [10]. In addition, treatment with selective serotonin reuptake inhibitors may lead to a decrease in platelet activity [11].

Mean platelet volume (MPV), which is indicative of platelet size, is accepted as an indication of platelet activity [12]. Increased MPV levels are thought to be closely related to cardiovascular disorders [13]. MPV values were studied in also schizophrenia, bipolar disorder and panic disorder with contradictory and inconclusive results [14–18]. We could not find any study evaluating the MPV levels in GAD patients in the literature. In this study we aimed to evaluate MPV values in GAD patients and explore the diagnostic value of MPV in GAD.

CONTACT Behice Han Almis  dr.behice@hotmail.com

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Method

Study population

In this retrospective cohort study, complete blood count especially in terms of platelet count (PLT), platelet distribution width (PDW), plateletcrit (PCT) and MPV values of 60 patients without any psychiatric treatment history who admitted to the psychiatry outpatient clinic of Adiyaman University Training and Research Hospital and diagnosed with GAD were compared with the data of 60 healthy subjects who were similarly distributed in regard to age (18–60 years) and gender. The diagnosis of GAD was made according to the fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) by an experienced psychiatrist.

Patients and controls whose data were missing or incomplete on record system were not included in the study. People with risk factors (drug use, hypertension, diabetes, coronary artery disease, cancer, hyperlipidemia, alcohol and substance use, pregnancy) that may have an impact on platelet counts were not included. Patients with other psychiatric comorbidities including unipolar depressive disorder were not included in the study even though their primary diagnosis were GAD.

This study was approved by the Adiyaman University Training and Research Hospital ethics committee (20.04.2017/3-10).

Biochemical analysis

Venous blood samples were obtained from antecubital vein of both patient and control group between 8 and 10 a.m. after at least 8 h of starving. Samples were centrifuged in 1 h and serum samples were studied at the same day in the biochemistry laboratory of Adiyaman University Training and Research Hospital with CELL-DYN 3700 SL analyzer (Abbott Diagnostics, Chicago, U.S.A.). The reference intervals were accepted as $142\text{--}424 \times 10^3/\mu\text{L}$ for platelet count, $0\text{--}1000$ fL for PDW, $0\text{--}1000\%$ for PCT and $6.8\text{--}10.8$ fL for MPV.

Statistical analysis

Statistical Package for Social Sciences (SPSS v20, Chicago, IL, U.S.A.) program for IOS were used. Descriptive parameters are expressed as mean, standard deviation or percentage. Kolmogorov–Smirnov test was applied to evaluate the normal distribution of the variables. Two-sided *t*-test and Pearson Chi-square test were used for comparison of normally distributed variables. Abnormal distributed variables were compared with Mann–Whitney *U*-test. Receiver operating characteristic (ROC) curve analysis was used to evaluate the diagnostic value of MPV. A value of less than .05 was considered statistically significant.

Results

The mean age was 39.82 ± 18.97 in the GAD group and 34.92 ± 14.60 in the control group. The ratio of females was 63.3% in the GAD group and 71.6% in the control group, and there was no significant difference in terms of age and gender between the two groups ($p = .229$ and $p = .330$, respectively).

According to the comparison of complete blood count values, MPV was significantly higher ($p = .008$) and platelet count was significantly lower in the GAD group ($p = .001$). Other variables showed no significant differences. A comparison of socio-demographic variables and complete blood count values of the patient and control groups are given in Table 1.

When the groups were separated according to the gender, MPV was significantly higher ($p = .010$) and platelet count was significantly lower in the women GAD group ($p = .001$) whereas there was no significant difference between control and GAD groups among men. Comparison of MPV and platelet counts of men and women are given in Table 2.

Correlation between age and complete blood count values also are given in Table 3 separately.

ROC curve analysis performed to assess the diagnostic value of MPV is shown in Figure 1. The area

Table 1. Comparison of socio-demographic variables and complete blood count values of patient and control groups.

		GAD	Control	<i>p</i>
Age (years)		38.35 ± 18.97	34.90 ± 14.62	.229
Gender	Women (n:81)	38 (%63.3)	43 (%71.6)	.330
	Men (n:39)	22 (%36.7)	17 (%28.4)	
WBC ($10^3/\mu\text{L}$)		8.04 ± 1.70	7.88 ± 1.84	.623
HGB (g/dL)		14.01 ± 1.42	13.64 ± 1.54	.177
HTC (%)		42.12 ± 3.98	41.74 ± 5.02	.644
RBC ($10^6/\mu\text{L}$)		4.86 ± 0.46	4.80 ± 0.55	.478
MCV (fL)		85.92 ± 4.93	85.84 ± 5.22	.936
PLT ($10^3/\mu\text{L}$)		258.83 ± 52.41	297.93 ± 57.85	.001*
PDW (fL)		19.44 ± 2.10	19.33 ± 1.25	.721
PCT (%)		0.21 ± 0.05	0.20 ± 0.04	.112
MPV (fL)		8.15 ± 1.41	7.50 ± 1.24	.008*

Notes: GAD: generalized anxiety disorder; WBC: white blood cell; HGB: haemoglobin; HTC: haematocrit; RBC: red blood cell; MCV: mean corpuscular volume; PLT: platelet; PDW: platelet distribution width; PCT: plateletcrit; MPV: mean platelet volume.

* $p < .05$.

Table 2. Comparison of MPV and platelet counts of men and women.

		GAD	Control	<i>p</i>
MPV (fL)	Women (n:81)	8.15 ± 1.43	7.38 ± 1.20	.010*
	Men (n:39)	8.15 ± 1.41	7.80 ± 1.33	.439
PLT (10 ³ /uL)	Women (n:81)	260.98 ± 53.52	308.13 ± 55.68	.001*
	Men (n:39)	255.11 ± 51.47	272.14 ± 56.72	.334

Notes: GAD: generalized anxiety disorder; PLT: platelet; MPV: mean platelet volume.

**P* < .05.

Table 3. The correlation between age and platelet parameters.

Platelet parameters	<i>r</i>	<i>p</i>
Red blood cell count	−0.075	.417
Haemoglobin	−0.091	.323
White blood cell count	−0.085	.358
Haematocrit	0.031	.735
Mean corpuscular volume	0.083	.367
Platelet count	−0.189	.039*
Platelet distribution Width	−0.104	.258
Plateletcrit	−0.311	.001**
Mean platelet volume	−0.228	.012*


**p* < .05.

***p* < .01.

under the ROC curve (AUC) of MPV levels for GAD was 0.655 (95% CI: 0.557–0.754, *p* < .001). The optimal cut-off value for MPV level was 7.45 fL, and its sensitivity and specificity for diagnosis of GAD were 65% and 56.7%, respectively.

Discussion

In recent years, studies have increasingly been made on platelets to understand some psychiatric disorders and to evaluate the pharmacological properties of psychiatric drugs. Platelets contain many clotting and growth factors, as well as serotonin (5-HT), ADP, ATP and calcium at high concentrations. They have α-2, β-2 adrenoceptors, benzodiazepine and 5-HT (1, 2, 6, 3) receptors on their membranes [19]. Markovitz et al. [20] found that emotional stress increases platelet activity and suggested that stressors may play an important role in vascular events such as myocardial infarction. In the case of increased stress and anxiety, serotonin binds to 5-HT-2 receptors on platelets and mediates the release of factors that promote platelet aggregation [21]. Pecknold et al. [22] found that platelet 5-HT reuptake was increased in patients with major depression accompanying panic disorder. With the guidance of psychopharmacological studies on platelet receptors, researchers have suggested that human platelet 5-HT-2A receptor is a valid peripheral indicator of brain 5-HT-2A receptors [23]. Moreover Garcia-Sevilla et al. [24] have found an increase in the number of platelet α-2 adrenergic receptors in mood disorders and suggested that platelet α-2 receptors may also be an indicator of brain α-2 receptors. In another study, subchronic psychological stress has

**Figure 1.** ROC curve analysis performed to assess the diagnostic value of MPV.

been shown to increase platelet α -2 receptor concentration, especially in women [25].

MPV has long been considered as an indicator of platelet activity [12]. The number of studies about MPV values in the field of psychiatry is scarce. In their large scale study, Canan et al. [26] showed that MPV values were increased in major depression patients, and these values were decreased after 8 weeks of escitalopram treatment. Results of the studies about MPV values in panic disorder patients were contradictory. Some reported that MPV values were higher in panic disorder patients [15,16] whereas others reported otherwise [14,17]. Increased MPV was attributed to activation of platelets via α -2 receptors and decreased MPV was attributed to 5-HT transporter rate and its metabolism. Serotonin imbalance, increased catecholamine levels and sympathetic system activation also play an important role in the pathogenesis of GAD and recent studies showed that these factors have important effects on platelet activation. Therefore increased MPV levels in GAD patients in our study supports the hypothesis of increased platelet activation due to sympathetic system activation. Also platelet levels were found to be significantly lower in GAD patients in our study. This finding is consistent with the non-linear inverse relation between platelet volume and platelet count [27].

Larger platelets inhabits more granules therefore have more vasoactive and prothrombic factors [28]. In a 10-year follow-up study published recently, GAD was found to be associated with cardiovascular mortality independent of metabolic syndrome and other cardiovascular risk factors. However, this relationship was detected only in female patients and not in male patients [29]. Supporting this result, when men and women were separately compared between patient and control groups in our study, only women GAD patients were found to have increased MPV levels. Even so, one should kept in mind that this finding may be due to small number of men patients and controls ($n = 39$). Nevertheless, it can be suggested that higher MPV levels especially in females may have a predictive diagnostic value for GAD. However, ROC curve analysis for differentiation of GAD from those in the control group demonstrated that MPV, at a cut-off value of 7.45 fL yielded a sensitivity and specificity of 56.44% and 87.72% respectively ($p < .001$). Unfortunately there is no established diagnostic marker for any psychiatric disorder yet and MPV too would be considered to be “poor” at separating GAD from healthy subjects.

Relationship between disease severity and MPV could not be determined since the study was performed retrospectively and there were no measurements of any severity scale was available in the records. The participants' body mass index and smoking status could not

be reached since the study was performed retrospectively. This is one of the important limitations. Another limitation of this study is the lack of data about specific inflammatory mediators which may have important effects on MPV values. The other limitation of this study is that power analysis was not used in the calculation of sample size. Despite all the limitations, this is the first study to evaluate MPV values in patients with GAD. We believe that further controlled prospective studies about this issue will be valuable.

Disclosure statement

No potential conflict of interest was reported by the authors.

ORCID

Ihsan Aksoy  <http://orcid.org/0000-0001-8426-2072>

References

- [1] Moffitt TE, Caspi A, Taylor A, et al. How common are common mental disorders? Evidence that lifetime prevalence rates are doubled by prospective versus retrospective ascertainment. *Psychol Med*. 2010;40(6):899–909.
- [2] Stein MB, Sareen J. Clinical practice. Generalized anxiety disorder. *N Engl J Med*. 2015;373(21):2059–2068.
- [3] Roest AM, Martens EJ, de Jonge P, et al. Anxiety and risk of incident coronary heart disease: a meta-analysis. *J Am Coll Cardiol*. 2010;56(1):38–46.
- [4] Piletz JE, Zhu H, Madakasira S, et al. Elevated P-selectin on trombosit in depression: response to bupropion. *J Psychiatr Res*. 2000;34(6):397–404.
- [5] Schins A, Hamulyák K, Scharpé S, et al. Whole blood serotonin and trombosit activation in depressed post-myocardial infarction patients. *Life Sci*. 2004;76(6):637–650.
- [6] Zafar MU, Paz-Yepes M, Shimbo D, et al. Anxiety is a better predictor of trombosit reactivity in coronary artery disease patients than depression. *Eur Heart J*. 2010;31(13):1573–1582.
- [7] Durmaz T, Özdemir Ö, Keleş T, et al. Trombosit and sympathetic activity in MI. *Turk J Med Sci*. 2009;39(2):259–265.
- [8] Thompson CB, Eaton KA, Princiotto SM, et al. Size dependent trombosit subpopulations: relationship of trombosit volume to ultrastructure, enzymatic activity, and function. *Br J Haematol*. 1982;50(3):509–519.
- [9] Vanhoutte PM. Trombosit-derived serotonin, the endothelium, and cardiovascular disease. *J Cardiovasc Pharmacol*. 1991; 17 (Suppl 5):6–12.
- [10] Parakh K, Sakhuja A, Bhat U, et al. Trombosit function in patients with depression. *South Med J*. 2008;101(6):612–617.
- [11] Galan AM, Lopez-Vilchez I, Diaz-Ricart M, et al. Serotonergic mechanisms enhance trombosit-mediated thrombogenicity. *Thromb Haemost*. 2009;102(3):511–519.
- [12] Tsiara S, Elisaf M, Jagroop IA, et al. Trombosit as predictors of vascular risk: is there a practical index

- of trombosit activity? *Clin Appl Thrombosis-Hemostasis*. 2003;9:177e90.
- [13] Slavka G, Perkmann T, Haslacher H, et al. Mean trombosit volume may represent a predictive parameter for overall vascular mortality and ischemic heart disease. *Arterioscler Thromb Vasc Biol*. 2011;31(5):1215–1218.
 - [14] Gül I G, Eryılmaz G, Ozten E, et al. Decreased mean trombosit volume in panic disorder. *Neuropsychiatr Dis Treat*. 2014;10:1665–1669.
 - [15] Asoglu M, Aslan M, Imre O, et al. Mean trombosit volume and red cell distribution width levels in initial evaluation of panic disorder. *Neuropsychiatr Dis Treat*. 2016;12:2435–2438.
 - [16] Kokacya MH, Copoglu US, Kivrak Y, et al. Increased mean trombosit volume in patients with panic disorder. *Neuropsychiatr Dis Treat*. 2015;11:2629–2633.
 - [17] Ransing RS, Patil B, Grigo O. Mean platelet volume and platelet distribution width level in patients with panic disorder. *J Neurosci Rural Pract*. 2017;8(2):174–178.
 - [18] Wysokinski A, Szczepocka E. Platelet parameters (PLT, MPV, P-LCR) in patients with schizophrenia, unipolar depression and bipolar disorder. *Psychiatry Res*. 2016;237:238–245.
 - [19] Camacho A, Dimsdale JE. Trombosits and psychiatry: lessons learned from old and new studies. *Psychosom Med*. 2000;62(3):326–336.
 - [20] Markovitz JH, Matthews KA. Trombosits and coronary heart disease: potential psychophysiological mechanisms. *Psychosom Med*. 1991;53:643–668.
 - [21] Garvey MJ, Noyes R, Woodman C, et al. The association of urinary 5-hydroxyindoleacetic acid and vanillylmandelic acid in patients with generalized anxiety. *Neuropsychobiology*. 1995;31(1):6–9.
 - [22] Pecknold JC, Suranyi-Cadotte B, Chang H, et al. Serotonin uptake in panic disorder and agoraphobia. *Neuropsychopharmacology*. 1988;1:173–176.
 - [23] Geaney DP, Schachter M, Elliot JM, et al. Characterization of [3H]lysergic acid diethylamide binding to a 5-hydroxytryptamine receptor on human trombosit membranes. *Eur J Pharmacol*. 1984;97:87–93.
 - [24] Garcia-Sevilla JA, Zis AP, Hollingsworth PJ, et al. Trombosit alpha-2 adrenergic receptors in major depressive disorder: binding of tritiated clonidine before and after tricyclic antidepressant drug treatment. *Arch Gen Psychiatr*. 1981;38:1327–1333.
 - [25] Maes M, Van Gastel A, Delmeire L, et al. Trombosit alpha2-adrenoceptor density in humans: relationships to stress-induced anxiety, psychasthenic constitution, gender and stress-induced changes in the inflammatory response system. *Psychol Med*. 2002;32(5):919–928.
 - [26] Canan F, Dikici S, Kutlucan A, et al. Association of mean trombosit volume with DSM-IV major depression in a large community-based population: the MELEN study. *J Psychiatr Res*. 2012;46(3):298–302.
 - [27] Bessman JD, Williams LJ, Gilmer PR Jr. Mean platelet volume. The inverse relation of platelet size and count in normal subjects, and an artifact of other particles. *Am J Clin Pathol*. 1981;76(3):289–293.
 - [28] Bath PM, Butterworth RJ. Trombosit size: measurement, physiology and vascular disease. *Blood Coagul Fibrinolysis*. 1996;7:157–161.
 - [29] Bunevicius A, Saudargiene A, Nemeroff CB, et al. Metabolic syndrome, major depression, generalized anxiety disorder, and ten-year all-cause and cardiovascular mortality in middle aged and elderly patients. *Int J Cardiol*. 2015;190:360–366.